



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

Problems 189 and 190 were also solved by L. E. Newcomb, Los Gatos, California.

190. Proposed by A. H. HOLMES, Brunswick, Maine.

Find the general term of the series 2, 3, 7, 46, 2112, etc.

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va., and GRACE M. BAREIS, Miss Roney's School, Bala, Pa.

Denoting the $n+1$ st term by T_{n+1} , we have the recursion formula $T_{n+1}=T_n^2-n$.

191. Proposed by NELSON L. RORAY, Bridgeton, N. J.

Find a number such that if it be multiplied by 2, 3, 4, 5, and 6, the cyclical order of its digits will not be changed.

Remark by L. E. DICKSON, Ph. D., The University of Chicago.

The answer is the period 142857, of the fraction $\frac{1}{7}$. Solved in the MONTHLY of 1895, p. 13, Algebra problem number 32.

GEOMETRY.

Problem 212 was also solved by F. D. Posey, A. B., San Mateo, California.

Problem 213 was also solved by L. E. Newcomb, Los Gatos, California, and F. D. Posey, A. B., San Mateo, California.

215. Proposed by M. J. NEWELL, A. M., Evanston High School, Evanston, Ill.

No satisfactory solution has been received.

216. Proposed by JOHN J. QUINN, Warren High School, Warren, Pa.

Find, by plane geometry, the sides of a right triangle if the hypotenuse is 35, and the side of the inscribed square is 12.

Solution by L. E. DICKSON, Ph. D., The University of Chicago.

“The inscribed square” to a right triangle is open to two interpretations: (1) the square has a side lying along the hypotenuse; (2) the square has two sides lying along the legs of the right triangle. For (1), the present problem is impossible (see problem 221). For the interpretation (2), denote the legs by a and b , $a \geq b$, whence $a^2 + b^2 = 35^2$. By similar triangles, $a - 12 : 12 = a : b$, whence $ab = 12(a + b)$. Adding and subtracting twice the latter from $a^2 + b^2 = 1225$, we get

$$(a+b)^2 - 24(a+b) = 1225, \quad (a-b)^2 = 1225 - 24(a+b).$$